AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

1.-10. (Canceled).

11. (Currently Amended) A process for the preparation of a metal-organic compound, comprising contacting at least one imine ligand according to formula 1, wherein the imine ligand is contacted with a metal-organic reagent of formula 2 in the presence of at least 1 equivalent of an inorganic or metal-organic base, or wherein an HA adduct of the imine ligand is contacted with a metal-organic reagent of formula 2 in the presence of at least 2 equivalents of an inorganic or metal-organic base, wherein formula 1 is:

Y=N-R

wherein Y is selected from a substituted carbon, nitrogen, or phosphorous atom and R represents a proton, a protic or an aprotic substitutent, and formula 2 is:

$$M^{V}(L_{1})_{k}(L_{2})_{l}(L_{3})_{m}(L_{4})_{n}X$$

wherein:

M represents a group 4 or group 5 metal ion,

V represents the valency of the metal ion, which is 3, 4 or 5,

L₁, L₂, L₃, and L₄ represent ligands on M which may be the same or different,

X represents a group 17-halogen atom,

k, l, m, n = 0, 1, 2, 3, 4 with k+l+m+n+1=V, and

wherein HA represents an acid, of which H represents its proton and A its conjugated base.

- 12. (Currently Amended) A process according to claim 11 wherein R represents a hydrogen atom and wherein-Y is selected from the group consisting of:
 - (i) a phosphorus substituent defined by the formula:

$$R^{11}$$
 R^{-12}
 R^{13} (formula 3)

wherein each R^{1j} , with j=1-3, is independently selected from the group consisting of a hydrogen atom, a halogen atom, a C_{1-8} alkoxy radical, a C_{6-10} aryl or aryloxy radical, an amido radical, or a C_{1-20} hydrocarbyl radical unsubstituted or substituted by a halogen atom, a C_{1-8} alkoxy radical, a C_{6-10} aryl or aryloxy radical, or an amido radical, or a silyl radical of the formula:

$$R^{21}$$
 $-Si-R^{22}$
 R^{23} (formula 4)

 Θ and a germanyl radical of the formula:

wherein R^{2j} is independently selected from the group consisting of hydrogen, a C_{1-8} alkyl or alkoxy radical or a C_{6-10} aryl or aryloxy radical radicals, provided each substituent R^{1j} or R^{2j} may be linked with another R^1 or R^2 to form a ring system, or and

(ii) a substituent defined by formula 6:



wherein each of Sub¹ and Sub² is independently selected from the group consisting of a hydrocarbyl radical radicals having from 1 to 30 carbon atoms; a-silyl radical radicals, a substituted or unsubstituted amido radical radicals and a-substituted or unsubstituted phosphido radical radicals, and wherein Sub¹ and Sub² may be linked with each other to form a ring system.

- 13. (Withdrawn/Currently Amended) A process according to claim 11, wherein the base is a carboxylate, or a fluoride, or a hydroxide, or a cyanide, or an amide, or a carbonate of Li, Na, K, Rb, Cs, or an ammonium salt, or a group 2 metal salt ehosenselected from Mg, Ca, or Ba, or an alkali metal ehosenselected from Li, Na, K, Rb, or Cs of a phosphate or a phosphate ester, and their aryl and alkyl compounds, or alkoxide and phenoxides of Li, Na, K, Rb or Cs or a phenoxide of Li, Na, K, Rb or Cs, or thallium hydroxide, alkylammonium hydroxide or a fluoride, or a hydrocarbanion of any of the group 1, group 2, group 12 or group 13 elements, or an alkali metal, group 1 hydride or group 2 hydride.
- 14. (Withdrawn) A process according to claim 13, wherein the inorganic base is selected from sodium hydride, or calciumhydride.

IJPEIJ et al Appl. No. 10/566,979 December 13, 2007

- 15. (Previously Presented) A process according to claim 11, wherein the metal-organic base is selected from an organolithium compound, or an organomagnesium compound.
- 16. (Previously Presented) A process according to claim 11, wherein the reaction is carried out in an aprotic solvent.
- 17. (Withdrawn) A process according claim 11, wherein the process is carried out in the presence of a phase transfer reagent.
- 18. (Previously Presented) A process for the preparation of a polyolefin by preparing a metal-organic compound according to the process of claim 11, wherein the base is an olefin polymerization-compatible base, wherein the metal-organic compound is activated anywhere in, or before a polymerization reactor.
- 19. (Previously Presented) A process according to claim 18, wherein the metal-organic compound is used without purification.
- 20. (Previously Presented) A process according to claim 18, wherein the metal-organic compound is formed in the polymerization reactor.